(22) International Filing Date:

#### WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5: (11) International Publication Number: WO 92/22283 A1 A61K 9/02, A61J 3/08 (43) International Publication Date: 23 December 1992 (23.12.92) (21) International Application Number: PCT/DK92/00187

(30) Priority data: 1165/91

17 June 1991 (17.06.91) DK

16 June 1992 (16.06.92)

(71) Applicant (for all designated States except US): FARMAC-EUTISK LABORATORIUM FERRING A/S [DK/ DKJ; Indertoften 5, DK-2720 Vanløse (DK).

(75) Inventor/Applicant (for US only): HALSKOV, Søren [DK/ DK]; Parcelvej 111, DK-2830 Virum (DK).

(74) Agent: HOFMAN-BANG & BOUTARD A/S; Adelgade 15, DK-1304 Copenhagen K (DK).

(81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CAPI p CH, CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (Eurk (European patent), GA (OAPI patent), GB, GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC (European patent), MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL, NL (European patent), NO, PL, RO, RU, SD, SE, SE (European patent), SN (OAPI patent), TD (OAPI patent), TG (OAPI patent), US.

#### Published

With international search report. With amended claims. In English translation (filed in Danish).

(54) Title: A PROCESS FOR PRODUCING SUPPOSITORIES BY COMPRESSION AND SUPPOSITORIES OBTAINED BY THE PROCESS

#### (57) Abstract

Suppositories which may have a considerably higher content of an active drug than ordinary suppositories, are produced by compressing a suppository mass, which contains a considerable amount of a polyethylene glycol having a molecular weight of at least 4000, preferably 6000. The polyethylene glycol constitutes 20-50 % by weight of the suppository mass, which moreover contains microcrystalline cellulose and/or other additives common in the production of drugs. The obtained suppositories are more convenient to store and use, easier to produce and considerably more concentrated with respect to active substance (up to 75 % by weight) than traditional melt-moulded suppositories.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT AU BB BE BF BG BI CA CF CG CH CM CO DE DIK ES	Austria Australia Barbados Belgium Berkina Faso Bolgaria Benin Brazil Canada Cantral African Republic Congo Switzer land C'ôte d'Ivoire Cameroon Czechoslovakia Germany Denmark Spain	FI FR GA GB GN GR HU IE IT JP KP LL LL MC MC	Finland France Gabon United Kingdom Goinea Greece Hungary Ireland Japan Democratic People's Republic of Korea Republic of Korea Linchtenstein Sri Lanka Luxembourg Monaco Madagascar	MI. MN MR MW NI. NO PL RO RU SD SE SN SU TD TG US	Mali Mongolia Mauritania Malawi Nutherlands Norway Poland Romania Russian Federation Sudan Sweden Senegal Soviet Union Chad Togo United States of America

A process for producing suppositories by compression and suppositories obtained by the process

\_\_\_\_\_

5

10

15

20

25

30

35

The present invention concerns a special process for producing suppositories by compression, and the process of the invention is characterized by producing a suppository mixture granulate containing a considerably greater amount of active drug than ordinary suppositories as well as 20-50% by weight of a polyethylene glycol having an average molecular weight of at least 4000, and producing the suppositories in the same manner as tablets, i.e. by compression, instead of by moulding. The invention moreover concerns the suppositories obtained by the process which have a considerably higher percentual content of active drug than ordinary suppositories.

As will be known, suppositories are drugs intended for insertion into the rectum. They contain the active drug in a dosed amount and are produced by pressing, moulding or compression. They can also be produced in the form of capsules for controlled release of the active substance.

Ordinarily suppositories are produced by moulding, the produced mass being melted using the least possible amount of heat, and then the liquid mass is poured into moulds having the desired nominal capacity.

The suppositories produced by moulding are oblong and smooth, and they have a uniform appearance. Melting is intended to provide a uniform distribution of the drug in the basic mass, which, however, can be difficult to obtain because of sedimentation during hardening.

However, traditional moulding is a time-consuming and slow process which involves considerable costs. Moulded suppo-

sitories also have the drawback that too strong heating of certain suppository basic masses result in unstable modifications with a considerably reduced solidification point.

5

It is well-known to use polyethylene glycols having average molecular weights of 4000-6000 or above as the main component in the basic mass for suppositories produced by traditional melting and moulding. Thus, the DE Offenle-gungsschrift 2 248 777 describes melt-moulded indomethacin suppositories whose basic mass contains such polyethylene glycols.

15

10

An object of the invention is to provide a process enabling the production of suppositories having a high content of active drug, which are partly more convenient to store and are partly more convenieny to use and easier and cheaper to produce than traditional melt-moulded suppositories.

20

This is achieved by the process of the invention which is characterized by the subject-matter stated in the characterizing portion of claim 1.

25

It has previously been attempted to produce suppositories by compression or pressing, i.e. by traditional tabletting methods. However, these suppositories tend to form irregular rough surfaces, which makes them unpleasant to use for the patient. Moreover, in such a production method it has been found impossible to dose the drug in so high doses as is often desirable owing to the prescribed treatment.

30

Thus, the EP publication 111 137 describes suppositories containing the drug indomethacin in a base consisting of polyethylene glycol having an average molecular weight of

10

15

20

25

30

35

up to 35,000. It is stated that the content of the active drug may be up to 50% by weight, but preferably the content is 2-40% by weight and in particular 2-26% by weight. All the examples in the publication concern rectal tablets having a content of indomethacin of 2.8 - 5.8% by weight and rectal capsules having a content of indomethacin of 5.25 - 10.5% by weight, i.e. rather low concentrations. Further, the suppositories thus known contain quite high amounts of polyethylene glycols, typically 1600-1730 mg per unit, which is a drawback, because it has been found that a content of polyethylene glycols in suppositories of 1 - 1.5 g per unit may cause bowel disorders.

It has now surprisingly been found that a composition which can easily be compressed to suppositories by an ordinary tabletting method can be obtained by using a suitable amount, more particularly 20-50% by weight of a polyethylene glycol having an average molecular weight of at least 4000 in the suppository basic mass. The use of such a polyethylene glycol in the basic mass results in suppositories having a uniform appearance and having an extremely smooth and regular surface, which is moreover sufficiently slippery for the suppository to be inserted without difficulty. It is moreover possible in such suppositories to incorporate up to 75% by weight of active drug, which is far above normal.

Polyethylene glycols, more particularly mixtures of condensation polymers of ethylene oxide and water, are also called "macrogols". Macrogols having average molecular weights of 200-700 are liquids, while macrogols having average molecular weights above 1000 vary in consistence from soft oily substances to hard wax-like solid substances. The average molecular weight is stated as a number after the name. "Macrogol 6000", which it is preferred to use according to the invention, thus has an ave-

10

15

20

25

rage molecular weight of about 6000.

The macrogols have the general formula

 $H(CH_2OCH_2)_nOH$ 

where n is greater than or equal to 4. In "Macrogol 6000" n has an average value of between 158 and 204. It is a white or cream-coloured solid wax-like substance which is in the form of a powder or flakes. The melting point is 56-63°C, i.e. considerably above the body temperature. Mixtures of various polyethylene glycols having melting points above the body temperature have frequently been used as a base in suppositories from which the drug is released by dissolution.

Since the melting points of the macrogols increase with increasing average molecular weight, macrogols having high average molecular weights, such as "Macrogol 6000", have not previously been a natural choice as a suppository base material.

Mixing a granulate of the drug with "Macrogol 6000" and compressing the mixture to suppositories by a method which is known per se for the production of tablets for oral administration provide the following advantages:

- The suppositories do not melt under normal temperature conditions, but only at about 60°C.
- 2) The suppositories are cheaper to produce.
  - 3) The drug can be incorporated in very high concentrations and is considerably easier to dose uniformly.

25

30

35

- 4) The product is more convenient and more pleasant to use for the patient.
- 5) Avoidance of drug decomposition during the heating which is necessary in traditional moulding of suppositionies.
- 6) It is easier to obtain and maintain a homogeneous mixture, and the risk of sedimentation is eliminated.

10 All active drugs which lend themselves for rectal administration may be incorporated in the suppositories produced according to the invention. One of the interesting drugs in this respect is 5-aminosalicylic acid (5-ASA), which is used particularly for the treatment of colitis ulcerosa 15 and Crohn's disease, but which has moreover been found to be of interest for the production of suppositories for treatment of hemorrhoids. Suppositories containing 5aminosalicylic acid are known e.g. from the EP patent application 83 775, these suppositories being produced by 20 moulding and containing max. 500 mg of 5-aminosalicylic acid per dose unit. The suppositories of the invention may contain considerably larger amounts of 5-ASA, which is absolutely an advantage for the patient.

An example of the many active drugs which may be used is steroids for various applications.

In addition to a polyethylene glycol or a mixture of polyethylene glycols having an average molecular weight of at least 4000, preferably "Macrogol 6000", in an amount of 20-50% by weight, the suppositories produced according to the invention contain microcrystalline cellulose and/or other additives common in the production of drugs. These additives and the active drug together amount to 45-75% based on the gross weight of the suppositories. Finally,

the suppositories contain one or more of the substances talc, magnesium stearate and polyvinyl pyrrolidone in an amount of 2-5% by weight.

The suppositories are typically compressed to symmetrical units having an approximately elliptic longitudinal section, i.e. the two ends are uniform (in contrast to the ordinary "torpedo-shape" where one end is pointed while the other is blunt).

The following example illustrates the invention:

#### **EXAMPLE**

A suppository basic mass for the production of 1000 suppositories consists of the following ingredients:

	"Macrogol 6000"	572 g
20	Microcrystalline cellulose	
	and active drug	1000 g
20	Magnesium stearate	4 g
	Talc	4 g
<i>y</i>	Polyvinyl pyrrolidone	
	+ ethanol (1:19)	q.s.

25

10

A granulate is made from the microcrystalline cellulose, the active drug, e.g. 5-aminosalicylic acid (5-ASA) and the mixture of polyvinyl pyrrolidone and ethanol.

The resulting granulate is mixed with "Macrogol 6000", and then magnesium stearate and talc are added.

The granulate can then be compressed to suppositories in a tabletting machine in a manner known per se.

10

15

20

## Patent Claims:

- 1. A process for producing suppositories by compression, c h a r a c t e r i z e d by producing a suppository mixture granulate containing a considerably larger amount of active drug than ordinary suppositories as well as 20-50% by weight of a polyethylene glycol having an average molecular weight of at least 4000, and then compressing the produced mass to suppositories by a method known per se for the production of tablets.
- 2. A process according to claim 1, c h a r a c t e r i z e d by producing a suppository mass having the following composition:

polyethylene glycol:

20-50왕

active drug

as well as microcrystalline cellulose

and/or other additives common in the

production of drugs:

45-75%, preferably

at least 50%

talc, magnesium stearate and/or

polyvinyl pyrrolidone:

2-5%

25

35

the stated percentages being percentages by weight.

- 3. A process according to claim 1 or 2, c h a r a c t e r i z e d in that the polyethylene glycol used has an average molecular weight of 6000.
  - 4. A process according to any of the preceding claims, c h a r a c t e r i z e d in that the polyethylene glycol used is selected from "Macrogol 4000", "Macrogol 6000" and mixtures thereof.

- 5. Suppositories, characterized in that they are produced by a process according to any of claims 1-4.
- 6. Suppositories according to claim 5, c h a r a c t e r i z e d in that they have the following composition:

polyethylene glycol having an average molecular weight of

average molecular weight of at least 4000:

20-50%

at least 4000:

active drug

as well as microcrystalline cellulose

and/or other additives common in the

production of drugs:

45-75%, preferably

at least 50%

talc, magnesium stearate and/or polyvinyl pyrrolidone:

2-5%

the stated percentages being percentages by weight.

20

- 7. Suppositories according to claim 5 or 6, c h a r a c t e r i z e d in that the polyethylene glycol used is "Macrogol 6000".
- 8. Suppositories according to any of claims 5-7, c h a r a c t e r i z e d in that they contain 5-aminosalicylic acid (5-ASA) as an active drug in an amount of up to 75% of the gross weight of the product.
- 9. Suppositories according to any of claims 5-7, c h a r a c t e r i z e d in that they contain a steroid as an active drug.

10

#### AMENDED CLAIMS

[received by the International Bureau on 23 October 1992 (23.10.92); original claims 1-9 replaced by amended claims 1-8 (2 pages)]

- 1. A process for producing suppositories by compression, c h a r a c t e r i z e d by producing a suppository mixture granulate containing 50-75% active drug as well as optionally microcrystalline cellulose and/or other additives common in the production of drugs, 2-5% talc, magnesium stearate and/or polyvinyl pyrrolidone as well as 20-50% by weight of a polyethylene glycol having an average molecular weight of at least 4000, and compressing the produced mass to suppositories by a method known per se for the production of tablets.
- 2. A process according to claim 1, c h a r a c t e r i z e d in that the polyethylene glycol used has an average molecular weight of 6000.
- 3. A process according to claim 1 or 2, c h a r a c 
  20 terized in that the polyethylene glycol used is
  selected from "Macrogol 4000", "Macrogol 6000" and mixtures thereof.
- 4. Suppositories, c h a r a c t e r i z e d in that they are produced by a process according to any of claims 1-3.
  - 5. Suppositories according to claim 4, c h a r a c t e r i z e d in that they have the following composition:

30

polyethylene glycol having an average molecular weight of at least 4000: active drug

20-50%

as well as microcrystalline cellulose and/or other

additives common in the production of drugs: 50-75% talc, magnesium stearate and/or polyvinyl pyrrolidone: 2-5%

5

10

15

the stated percentages being percentages by weight.

- 6. Suppositories according to claim 4 or 5, c h a r a c t e r i z e d in that the polyethylene glycol used is "Macrogol 6000".
- 7. Suppositories according to any of claims 4-6, c h a r a c t e r i z e d in that they contain 5-aminosalicylic acid (5-ASA) as an active drug in an amount of up to 75% of the gross weight of the product.
- 8. Suppositories according to any of claims 4-6, c h a r a c t e r i z e d in that they contain a steroid as an active drug.

20

25

### INTERNATIONAL SEARCH REPORT.

International Application No. PCT/DK 92/00187

International Application No PCI/DR 32/0010/										
CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) <sup>8</sup> According to International Patent Classification (IPC) or to both National Classification and IPC										
IPC5: A 61 K 9/02, A 61 J 3/08										
II. FIELDS SEARCHED										
Minimum Documentation Searched 7										
Classificat	ian System	Classification Symbols								
'										
TDCS	A 61 W. A 61 1	·								
1703	IPC5 A 61 K; A 61 J									
Documentation Searched other than Minimum Documentation to the Extent that such Documents are included in Fields Searched <sup>2</sup>										
SE DK I	FI,NO classes as above									
	MENTS CONSIDERED TO BE RELEVANTS									
Category *	Citation of Document,11 with Indication, where a	· · · · · · · · · · · · · · · · · · ·	Relevant to Claim No. <sup>13</sup>							
X	DE, A, 2248777 (MERCK & CO. IN	C.)	1-9							
	19 April 1973,									
	see the whole document									
			,							
X	EP, A2, 0111137 (TROPONWERKE G	MBH & CO. KG)	1-9							
	20 June 1984,									
	see the whole document		·							
		·	·							
			İ							
	·	·								
- [			•							
ı		İ								
* Specia	categories of cited documents; 10	"T" (stor document outsighed effor t	he international filing date							
	ment defining the general state of the art which is no ildered to be of particular relevance	"T" later document published after to or priority date and not in confli- cited to understand the principle	or theory underlying the							
E Soli	ler document but published on or after the internations g date	invention								
		"X" document of particular relevance cannot be considered novel or ca involve an inventive step	innot be considered to							
whic	ment which may throw doubts on priority claim(s) or th is cited to establish the publication date of another ion or other special reason (as specified)	"Y" document of particular relevance	the claimed invention							
which is clied to establish the publication date of another citation or other special reason (as specified)  "O" document referring to an oral disclosure, use, exhibition or other means  "Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such document is combined with one or more other such document.										
"P" document published prior to the international filing date but international filing date but international filing date but "&" document member of the same patent family										
ister than the priority date claimed a document member of the same parcent rainty										
Date of the Actual Completion of the International Search   Date of Mailing of this International Search Report										
21st September 1992 <b>2</b> 2 -09- 1992										
International Searching Authority Signature of Authorized Officer										
huner tonson										
SWEDISH PATENT OFFICE Anneli Jönsson										
1000000	/210 (second sheet) (January 1986)									

# ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.PCT/DK 92/00187

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the Swedish Patent Office EDP file on The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)		Publication data
DE-A- 2248777	73-04-19	AT-B- AU-B- AU-D- BE-A- CA-A- FR-A-B- GB-A- NL-A- SE-B-C- US-A-	336188 472229 4708472 789726 985625 2158209 1384760 7212626 384134 3849549	77-04-25 76-05-20 74-04-04 73-04-05 76-03-16 73-06-15 75-02-19 73-04-10 76-04-26 74-11-19
EP-A2- 0111137	84-06-20	DE-A- JP-A-	3241263 59101420	84-05-10 84-06-12